

The rotation-vibration structure of ^{12}C

Gai, M.; Bijker, R.; Freer, M.; Kokalova Wheldon, Tzanka; Marin-Lambarri, D. J.; Wheldon, C.

DOI:

[10.1088/1742-6596/569/1/012011](https://doi.org/10.1088/1742-6596/569/1/012011)

License:

Creative Commons: Attribution (CC BY)

Document Version

Publisher's PDF, also known as Version of record

Citation for published version (Harvard):

Gai, M, Bijker, R, Freer, M, Kokalova Wheldon, T, Marin-Lambarri, DJ & Wheldon, C 2014, 'The rotation-vibration structure of ^{12}C ', *Journal of Physics: Conference Series*, vol. 569, no. 1, 012011.
<https://doi.org/10.1088/1742-6596/569/1/012011>

[Link to publication on Research at Birmingham portal](#)

General rights

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact UBIRA@lists.bham.ac.uk providing details and we will remove access to the work immediately and investigate.

The Rotation-Vibration Structure of ^{12}C

This content has been downloaded from IOPscience. Please scroll down to see the full text.

2014 J. Phys.: Conf. Ser. 569 012011

(<http://iopscience.iop.org/1742-6596/569/1/012011>)

View [the table of contents for this issue](#), or go to the [journal homepage](#) for more

Download details:

IP Address: 147.188.40.113

This content was downloaded on 23/12/2014 at 15:54

Please note that [terms and conditions apply](#).

The Rotation-Vibration Structure of ^{12}C

M Gai¹, R Bijker², M Freer³, T Kokalova³, D J Marin-Lambarri³, C Wheldon³

1. LNS at Avery Point, University of Connecticut, Groton, CT 06340-6097, USA
and WNSL, Dept of Physics, Yale University, New Haven, CT 06520-8124

2. Instituto de Ciencias Nucleares, Universidad Nacional
Autonoma de Mexico, A.P. 70-543, 04510 Mexico, D.F., Mexico

3. School of Physics and Astronomy, University of Birmingham, Birmingham B15 2TT, UK

E-mail: moshe.gai@yale.edu

Abstract. The newly measured high spin $J^\pi = 5^-$ state at 22.4(2) MeV in ^{12}C reported in this conference, fits very well to the predicted (ground state) rotational band of an oblate equilateral triangular spinning top with a \mathcal{D}_{3h} symmetry characterized by the sequence of states: $0^+, 2^+, 3^-, 4^\pm, 5^-$ with almost degenerate 4^+ and 4^- (parity doublet) states. Such a \mathcal{D}_{3h} symmetry was observed in triatomic molecules, and it is observed here for the first time in nuclear physics. We discuss a classification of other rotation-vibration bands in ^{12}C such as the (0^+) Hoyle band and the (1^-) bending mode band and suggest measurements in search of the predicted (“missing”) states that may shed new light on clustering in ^{12}C and light nuclei. In particular, the observation (or non observation) of the predicted (“missing”) states in the Hoyle band will allow us to conclude the geometrical arrangement of the three alpha particles composing the Hoyle state at 7.654 MeV in ^{12}C .

1. Introduction

The recent observation of the 2^+ Hoyle rotational excitation in ^{12}C [1] allows for the first time the study of the Rotation-Vibration Structure of ^{12}C . These new data and the recently discovered 4^- [2, 3] and 4^+ [4] states in ^{12}C are in agreement with the predicted spectrum of an oblate spinning top with a \mathcal{D}_{3h} symmetry [5, 6]. It was predicted [5, 6] that the three alpha-particle system of ^{12}C leads to the ground state rotational band including the most unusual sequence of states: $0^+, 2^+, 3^-, 4^\pm, 5^-$. The new high spin 5^- state reported by the Birmingham group in this conference [7] as well the previously published 4^- state [2, 3] lead to a $J(J+1)$ trajectory as predicted by this U(7) model [8, 9] including the nearly degenerated 4^- and 4^+ states as shown in Fig. 1.

2. The Algebraic Cluster Model

The spectrum of ^{12}C predicted by the Algebraic Cluster Model (ACM) [5, 6] is shown in Fig. 2 where it is also compared to the measured spectrum of ^{12}C [8]. In addition to the ground state rotational band this U(7) model [5, 6] predicts the Hoyle state at 7.65 MeV in ^{12}C to be the first vibrational breathing mode of the three alpha-particle equilateral configuration leading to the same rotational structure albeit with a larger moment of inertia (by a factor of 2). Recent measurements revealed the 2^+ [1] and 4^+ [4] members of the predicted Hoyle rotational band and we are currently searching [10] for the 4^- predicted by the ACM to be nearly degenerated






Figure 1. Rotational band structure of the: ground-state band, the Hoyle band and the bending vibration band in ^{12}C .




Figure 2. Comparison between the low-lying experimental spectrum of ^{12}C and the energies of the oblate symmetric top. The levels are organized in columns corresponding to the ground state band and the vibrational bands with A and E symmetry [5, 6] of an oblate top with triangular symmetry. The last column on the left-hand side, shows the lowest observed non-cluster (1^+) levels.

with the 4^+ state and the 3^- (broad) state that was suggested to lie between 11 and 14 MeV [2]. The observation (or lack there) of these “missing” states will allow us to determine whether the Hoyle state is composed of three alpha-particles in an equilateral triangle arrangement [8, 11, 12] or an obtuse triangle [13] or whether it is better described as vibrational excitation of a “diffuse gas” of three alpha-particles [14].

The $U(7)$ model also predicts the 1^- state at 10.84 MeV to be the vibrational bending mode with a rotational band including the 1^- and a degenerate 2^\pm states. We are searching [10] for the third 2^+ of ^{12}C that is predicted by the $U(7)$ model to lie near the observed 2^- state at 11.8 MeV.

In conclusion the ACM appears to open a new chapter in cluster physics of light nuclei and it presents an opportunity for further experimental investigation of light nuclei.

3. Acknowledgments

This material is based upon work supported by the U.S. Department of Energy, Office of Science, Office of Nuclear Physics, under Award Number DE-FG02-94ER40870.

References

- [1] Itoh M *et al.* 2004 Nucl. Phys. **A738**, 268; Itoh M *et al.* 2011 Phys. Rev. C **84**, 054308; Freer M *et al.* 2009 Phys. Rev. C **80**, 041303(R); Zimmerman W R *et al.* 2011 Phys. Rev. C **84**, 027304(BR); Zimmerman W R *et al.* 2013 Phys. Rev. Lett. **110**, 152502
- [2] Freer M *et al.* 2007 Phys. Rev. C **76**, 034320 (2007)
- [3] Kirsebom O S *et al.* 2010 Phys. Rev. C **81**, 064313
- [4] Freer M *et al.* 2011 Phys. Rev. C **83**, 034314
- [5] Bijker B and Iachello F 2000 Phys. Rev. C **61**, 067305
- [6] Bijker R and Iachello F 2002 Ann. Phys. **298**, 334
- [7] Marin-Lambarri D J 2014 Invited talk, SOTANCP3, Yokohama, Japan, May 26-30
- [8] Marin-Lambarri D J, Bijker R, Freer M, Gai M, Kokalova Tz, C. Wheldon, 2014 Phys. Rev. Lett. **113**, 012502.
- [9] Popular press of Ref. [8]: Kevin Dusling, Physics, Synopsis:
<http://physics.aps.org/synopsis-for/10.1103/PhysRevLett.113.012502>,
 Hamish Johnston, Physicsworld.com: <http://physicsworld.com/cws/article/news/2014/jul/08/carbon-nu>.
- [10] A $^{12}\text{C}(e,e')$ experiment in progress at Darmstadt, P. von Neumann-Cosel, private communication, 2013.
- [11] Chernykh M, Feldmeier H, Neff T, von Neumann-Cosel P, and Richter A 2007 Phys. Rev. Lett. **98**, 032501
- [12] Yoshiko Kanada-Enyo 2007 Prog. of Theo. Phys. **117**, 655
- [13] Epelbaum E, Krebs H, Lee D, Meissner Ulf-G 2012 Phys. Rev. Lett. **106**, 192501; Epelbaum E, Krebs H, Lahde T, Lee D, and Meissner Ulf-G 2012 Phys. Rev. Lett. **109**, 252501
- [14] Funaki Y, Horiuchi H, von Oertzen W, Ropke G, Schuck P, Tohsaki A, and Yamada T 2009 Phys. Rev. C **80**, 64326 and references therein